

# The MONS Star Trackers<sup>1</sup>

Timothy R. Bedding

*School of Physics, University of Sydney 2006, Australia*

Hans Kjeldsen

*Teoretisk Astrofysik Center, Danmarks Grundforskningsfond, Aarhus University, DK-8000 Aarhus C, Denmark*

## ABSTRACT

The *MONS* satellite will have two Star Trackers to sense the spacecraft attitude, and we plan to use them as scientific instruments to perform high-precision photometry of thousands of stars. We briefly describe the current plans for the Star Trackers and their expected capabilities.

## 1. Introduction

Like many satellites that require precise attitude control, *MONS* will use star trackers to sense the spacecraft attitude. A star tracker is basically a wide-field CCD camera. In acquisition mode, images of the sky are compared with a star catalogue to determine the absolute orientation of the spacecraft. Once a target is acquired, the spacecraft attitude is continuously updated in tracking mode.

There is clear scientific interest in using the photometry from a star tracker to observe variable stars. Despite the small aperture, the advantages of space (long observing periods and no atmospheric scintillation) make such a camera superior to ground-based telescopes for some applications. The only example so far has been the work by Buzasi et al. (2000 and these Proceedings), who are making impressive use of the 52-mm star camera on the *WIRE* satellite after the failure of the main instrument. As far as we know, *MONS* is the first mission to be designed from the start to use star trackers for science. Of course, we must keep in mind that the primary role of the star trackers is to sense spacecraft attitude with the precision required for the main camera.

We describe here the plan for the *MONS* Star Trackers as it stood at the time of the Workshop (January 2000). This plan differs slightly from that described in the *MONS* Proposal (Kjeldsen, Bedding & Christensen-Dalsgaard 1999), and will undoubtedly change again before the final design is frozen<sup>2</sup>. There are several factors driving the changes, including: (i) performance of trackers as attitude sensors for the main camera, including redundancy issues; (ii) volume, thermal and power constraints from the possible presence of *Ballerina*; and (iii) performance of trackers as secondary science instruments.

---

<sup>1</sup>To appear in *Proceedings of the Third MONS Workshop: Science Preparation and Target Selection*, edited by T.C.V.S. Teixeira and T.R. Bedding (Aarhus: Aarhus Universitet).

<sup>2</sup>One change is that we now use the term “Star Tracker” rather than “Star Imager.”



Fig. 1.— The TERMA Star Tracker.

## 2. Current plans for the Star Trackers

There will be two Star Trackers, pointing in opposite directions. Both will consist of a small CCD camera (24 mm aperture) with a circular field of view with diameter  $22^\circ$  (see Fig. 1). The Bright Star Tracker (BST) will point forwards, in the same direction as the main telescope, and will have a broad blue filter (380–420 nm) to allow it to observe bright stars without saturating. The Faint Star Tracker will be unfiltered, preventing observations of very bright stars but improving the signal-to-noise at the faint end.

Figures 2–4 show the expected photometric performance of the two Star Trackers. As a guide, the FST will take several seconds to reach the same photometric precision for a given star as an individual *Hipparcos* measurement. Of course, each star was only measured by *Hipparcos* about 100 times over a few years, whereas *MONS* will observe each field almost continuously for about one month.

We intend to use the *Hipparcos* catalogue as the input catalogue for *MONS*. There will be about 200 stars in a typical BST field ( $V = -0.5$  to  $8.0$ ), rising to about 600 in the plane of the Milky Way. For the FST ( $V = 3.5$  to  $12.0$ ) the corresponding numbers are about 700 and 2000 stars.

The number of stars that will be measured and down-linked will depend both on crowding and down-link capacity. As a general rule one can assume that *MONS* BST will measure all *Hipparcos* stars down to  $V = 7$  in the non-Milky Way fields, except stars that are too crowded. For the FST, the equivalent value is  $V = 8.5$ . Stars below these magnitude limits will only be observed if they are of particular scientific interest.




Fig. 2.— The expected photometric precision per second for the two Star Trackers. The dashed parts of each curve indicate regions where the noise will probably be too high to give useful data. The diagonal dotted lines show the limits set by photon noise.




Fig. 3.— Expected  $3\sigma$  sensitivity to oscillations after 30 days for the two Star Trackers.




Fig. 4.— Expected noise levels after 30 days for the two Star Trackers, normalised by the solar oscillation signal in the two passbands. The BST benefits from the fact that oscillations are stronger in the blue than the red.

### 3. Possible changes to the plan

Details subject to change include the directions in which the Star Trackers point and the filters used. For example, one might consider having the two Star Trackers pointing in the same direction, one with a red filter and the other with a blue filter. This would give colour information, which is of great scientific value for many types of stars. The full range of observable magnitudes could be retained by using bracketed exposure times, with the CCD being read out in a sequence (e.g., 50 ms, 200 ms and 1000 ms) to sample both bright and faint stars. The total number of stars observed in this revised configuration would be similar to that described in Section 2, but in two colours and with the stars concentrated in half the number of fields.

### *Acknowledgements*

We thank the Australian Research Council for financial support, as well as the Danish National Research Foundation through its establishment of the Theoretical Astrophysics Center.

### REFERENCES

- Buzasi, D. L., Catanzarite, J., Laher, R., Conrow, T., Shupe, D., Gautier III, T. N., & Kreidl, T., 2000, *Astrophys. J., Lett.* (in press)

Kjeldsen, H., Bedding, T. R., & Christensen-Dalsgaard, J. (eds.), 1999, *MONS: Measuring Oscillations in Nearby Stars. A proposal for the Danish Small Satellite Programme*, Aarhus: Aarhus Universitet